




EGYPTIAN ACADEMIC JOURNAL OF
BIOLOGICAL SCIENCES
MICROBIOLOGY

G



ISSN
2090-0872

WWW.EAJBS.EG.NET

Vol. 16 No. 1 (2024)



Genetic Characterization of Three Soil *Streptomyces* Isolates Exhibiting Antiviral Activity Against TMV; Insights from 16S rRNA Gene Sequencing and RAPD Analysis

Behairy Akl¹ and Samar S. El-Masry²

¹Agricultural Microbiology Department, Faculty of Agriculture, Zagazig University, B.O. Pox 44519, Zagazig, Egypt.

²Dept. of Agric. Microbiol. (Lab. of Virology), Fac. Agric., Ain Shams University, P.O. Box 68, Hadayek Shobra 11241, Cairo, Egypt.

*E.Mail: samar_saved@agr.asu.edu.eg

ARTICLE INFO

Article History

Received:12/3/2024

Accepted:16/4//2024

Available:20/4/2024

Keywords:

Streptomyces,
Antiviral, TMV,
16SrRNA,
RAPD,
Biological
classification,
Identification.

ABSTRACT

In the present study, each of the 16S ribosomal RNA (rRNA) gene sequences and random amplified polymorphic DNA (RAPD) method were used to confirm the biological classification of three *Streptomyces* isolates having antiviral activities against Tobacco mosaic virus (TMV). The strongest anti-TMV effect with an inhibition percentage of 97.5 was recorded for *S. rochei*-IS-02 followed by *S. caeaoi*-IS-03 (95.5%) and *S. aureocirculatus*-IS-01 (92.5%) at dilution 1/2. Lengths of 771, 663 and 693 nts were determined for the partial nucleotide sequences of 16SrRNA gene of *S. aureocirculatus*-IS-01, *S. rochei*-IS-02 and *S. caeaoi*-IS-03, respectively. Sequences producing significant alignments showed identities ranging from 98.41 to 100%. The phylogenetic relationship confirmed the identification of the *Streptomyces* isolates. A total of 85 DNA fragments (81 polymorphic and 4 monomorphic) were amplified in RAPD assay of the three *Streptomyces* isolates using eight random primers and contained unique DNA fragments as positive DNA markers. These DNA fragments were distributed as 48, 41 and 42 for the three isolates. The eight primers produced a total amplified DNA fragments of 85 distributed as follows 8, 10, 13, 12, 12, 10, 11 and 9 for the eight oligonucleotide primers, respectively. The similarities between the three *Streptomyces* species based on RAPD-PCR ranged from 66.332 to 76.158%. The phylogenetic tree showed that *S. aureocirculatus*-IS-01 and *S. rochei*-IS-2 were found together in the same cluster, while, *S. caeaoi*-IS-03 was found in a separate cluster. As a conclusion 16SrRNA and RAPD-PCR methods were confirmed to be more than suitable tools to differentiate between the *Streptomyces* species.

INTRODUCTION

Streptomyces were successfully isolated, purified and characterized from different soils worldwide including Egypt (Mohamed *et al.*, 2001, Mohamed *et al.*, 2012, Shori *et al.*, 2012, Alvarez-Mico *et al.*, 2013, Kamjam *et al.*, 2019 and Bo *et al.*, 2019).

According to previous investigations, it was proved that the secondary metabolites extracted from actinomycetes were important in controlling plant diseases including plant viruses (Chen *et al.*, 2019). They also reported that the *Streptomyces* strain STZ isolated from the soil of Tianzhu Mountain in Shenyang, China exhibited antiviral activity against TMV. Belonging to 65 families 885 plant species were susceptible to infection of TMV (Roossinck 2015, Fraile and García-Arenal, 2018).

It is very difficult to rely on the morphological identification and cultural and physiological characteristics of actinobacteria including streptomycete (Kim *et al.*, 1999). This is due to the high similarity between them particularly in the color of aerial mycelium, type of spore chain, production of melanoid pigments and spore surface.

The use of molecular biology methods such as 16S ribosomal RNA (16SrRNA) gene could be considered an ideal tool to confirm biological identification (Anderson and Wellington 2001). 16SrRNA and 18SrRNA were applied to identify four bacterial (*Bacillus* sp. MSLB-1, *Bacillus* sp. MSLB2, *Bacillus* sp. MSLB3, *Escherichia coli* MSL-19) and three fungal isolates (*Aspergillus niger* MLSAs1, *Aspergillus* sp. MLSAs2, *Penicillium* sp. MLSP1), respectively, isolated from soil up to strains (El-Masry *et al.*, 2019).

16SrRNA was applied to identify four bacterial (*Bacillus* sp. MSLB-1, *Bacillus* sp. MSLB2, *Bacillus* sp. MSLB3, *Escherichia coli* MSL-19) isolated from soil up to strains (El-Masry *et al.*, 2019). Sequence data of 16SrRNA revealed that it could provide the basis for proposals to unify the genus *Streptomyces*, in addition to other phylogenetic investigations that sought to determine the intrageneric structure of a genus (Stackebrandt *et al.*,

1991, Mehling *et al.*, 1995, Kim *et al.*, 1996 and Takeuchi *et al.*, 1996).

Biswas *et al.* (2017) used 16S ribosomal RNA (rRNA) gene sequence to confirm the characterization and identification of some streptomycetes (*Streptomyces labedae*, *S. variabilis*, *S. erythrogriseus* and *S. griseoincarnatus*) isolated from sediment of Sundarbans mangrove forest in India, previously classified based on medium 2 in International *Streptomyces* Project (ISP), the isolates characterized by a yellowish brown to red aerial hyphae, retinaculum-apertum chain spore with spiny-surface.

El-Masry *et al.* (2021) isolated, purified and identified some bioagent streptomycetes having antimicrobial activities from the soil in Egypt followed by molecular identification up to strains based on 16SrRNA gene.

Physiological characteristics represented using sugars as a sole carbon source and the production of enzymes cannot be relied upon absolutely in the identification of such microorganisms. Therefore, it is necessary to use the recent techniques that can differentiate between them such as random amplified polymorphic DNA (RAPD) (Malkawi *et al.*, 1999, Gharaibeh *et al.*, 2003, Hussein *et al.*, 2006), inter simple sequence repeat (ISSR) (Bornet and Branchard 2001, Bornet *et al.*, 2002, Fedrigo *et al.*, 2016 and Abd El-Aziz *et al.*, 2023) among DNA fingerprinting of streptomycetes.

RAPD-PCR technique was applied in several investigations to differentiate between streptomycetes isolated from different sources, *i.e.*, soil and marine ecosystems (Hussein *et al.*, 2006, Luo *et al.*, 2010, Mohamed *et al.*, 2012, Shori *et al.*, 2012, Mohamed, *et al.*, 2013, Maleki *et al.* 2013, Biswas *et al.* 2017, Qattan and Khattab 2019 and Shoukry *et al.* 2019).

Malkawi *et al.* (1999) concluded that RAPD-PCR method was efficient in discriminating and evaluating the genetic

variation between the *Streptomyces* isolates. Martin *et al.* (2000) evaluated six oligonucleotides presenting a high G+C content as RAPD primers were selected to compare 12 strains of six *Streptomyces* species. Therefore, RAPD-PCR technique allowed the rapid, sensitive and specific evaluation of genetic diversity among *Streptomyces* species.

This study was designed to use each of 16SrRNA gene and RAPD marker to confirm and differentiate between three soil-*Streptomyces* isolates having antiviral activities against TMV.

MATERIALS AND METHODS

Source of *Streptomyces* Isolates:

The three *Streptomyces* isolates under investigation were kindly provided by the Department of Agricultural Microbiology, Faculty of Agriculture, Ain Shams University, Cairo, Egypt. These isolates were biologically identified as *S. aureocirculatus*, *S. rochei* and *S. caeaoi*, and coded as IS-01, IS-02 and IS-03, respectively.

Preparation of *Streptomyces* Suspension:

Two spore discs of the three *Streptomyces* isolates were separately added to 250 mL flasks containing 50 mL of ISP3 broth medium as recommended by Chen *et al.* (2019) followed by incubation for a week at 28°C under shaking in a rotary shaker at 180 rpm. Low-speed centrifugation at 8000 rpm for 20 min was done, and the pellets were Two spore discs of the three *Streptomyces* isolates that were separately added to 250 mL flasks containing 50 mL of ISP3 broth medium as recommended by Chen *et al.* (2019) followed by incubation for a week at 28°C under shaking in a rotary shaker at 180 rpm. Low-speed centrifugation at 8000 rpm for 20 min was done, and the pellets were discarded, and the filtrates were stored at 4 °C till use.

Antiviral Activities of *Streptomyces* Strains against TMV:

An indicator host (*Nicotiana glutinosa*) inducing necrotic local lesions

(NLL) with TMV and containing 4-5 leaves was used. The infectious TMV sap was prepared according to the method of El-Ahdal *et al.* (1984). The crude TMV extract was diluted by adding an equal volume of 0.01 M phosphate-buffered saline (PBS), pH 7.0 before use followed by preparation of two-fold dilutions up to 1/16. Antiviral activities of the three streptomyces isolates were against TMV were determined against TMV using the diluted infectious sap of TMV by the half-leaf method as described (Gooding and Hebert 1967). The inoculated leaves were washed with sterile distilled water after 30 min from inoculation and cultivated in a glass greenhouse at 26 ±2°C. Four/five days post inoculation the number of NLL was recorded. The percentage of inhibition was calculated based on the following formula: 100- (Average number of NLL with diluted filtrate treatment ÷ No. of NLL at control leaf × 100).

16SrRNA Gene Sequencing and Analysis:

The three *Streptomyces* isolates which are biologically classified as *S. aureocirculatus*, *S. rochei* and *S. caeaoi* were sent to Macrogen® (908 world Meridian venture center, #60-24, Gasandong, Geumchun-gu, Seoul 153-781, Korea) for determining the nucleotide sequences of 16SrRNA gene for confirmation of the biological identification. Two universal primers called 785F: GGA TTA GAT ACC CTG GTA 3'; 907R: 5' CCG TCA ATT CMT TTR AGT TT3' were applied in the presence of DNA extracts of the three isolates which were prepared according to the method of Mohamed *et al.* (2012). Using the protocol supplied by the manufacturer, the nucleotide sequences were performed using the ABI PRISM BigDye™ Terminator Cycle Sequencing Kits, ABI PRISM 3730XL Analyzer (96 capillary type) sequencer (Applied Biosystems), MJ Research PTC-225 Peltier Thermal Cycler, DNA polymerase (FS enzyme, Applied Biosystems). The nucleotide sequences were aligned with the most similar strains documented in GenBank, using Standard Nucleotide

BLAST, National Library of Medicine, NCBI

(<https://blast.ncbi.nlm.nih.gov/Blast.cgi?PROGRAM>).

RAPD DNA Fingerprinting:

DNA extracts of the three *Streptomyces* isolates were prepared according to the method of Mohamed *et al.* (2012) and used as templates for PCR amplification in the presence of eight random oligonucleotides belonging to different groups: P-01 (5'AGG GGT CTT G3'), P-02 (5'CAATCGCCGT3'), P-03 (5'GTG AGG CGT C3'), P-04 (5'CCG CAT CTA C3'), P-05 (5'TGA GTG GGT G3'); P-06 (5'-CTT CCC CAA G-3'), P-07 (5'TCA GGG AGG T3'), P-08 (5'GGT GCA CGT T3'), following the PCR protocol described by Mohamed *et al.* (2013). DNA polymorphisms were scored as I (Present) and 0 (Absent), and each genetic similarity, pairwise combination and genetic distance were recorded (Hassanien *et al.* 2004) which was used to determine population relationships using the unweighted pair group method of analysis (UPGMA) (Sneath and Sokal 1973). On PCR the amplified products were visualized by electrophoresis (Sambrook *et al.*, 1989) in a 1.2% agarose gel at 60 v for 2.5 h with 1X TBE buffer. DNA fragments appeared in the gel after UV light transilluminator examination.

RESULTS AND DISCUSSION

In the present study, two *Streptomyces* isolates (*S. rochei*-IS-02, *S.*

caeaioi-IS-03) belonging to the grey series and one isolate belonging to the white series (*S. aureocirculatus*-IS-01) were used. They carry spore chains of RF and Spiral with smooth surfaces (Figure 1). *S. caeaioi*-IS-03 was producing Melanoid pigment and they varied in their ability to grow on Czapek's medium. Ten sugars (D-Fructose, D-Galactose, D-Glucose, D-Mannitol, D-Xylose, i-Inositol, L-Arabinose, L-Rhamnose, Raffinose and Sucrose) were used as a sole source of carbon by *S. caeaioi*-IS-03 in the growth medium. *S. aureocirculatus*-IS-01 and *S. rochei*-IS-02 can't utilize Raffinose. *S. rochei*-IS-02 was not able to utilize sucrose as a sole source of carbon. A similar was noted by *S. aureocirculatus*-IS-01 as it was not able to utilize L-Arabinose and L-Rhamnose, while *S. rochei*-IS-02 can't use Raffinose.

In 1943, Waksman and Henrici proposed the *Streptomyces* genus as a spore-forming, aerobic gram-positive bacteria. These bacteria induced extensive branching substrates and aerial mycelia and were characterized with DNA with a high content of cytosine and guanine which ranged from 69 to 73 1% (Williams *et al.*, 1983, Williams *et al.*, 1989 and Korn-Wendisch and Kutzner, 1992). This type of bacteria is also known for the presence of LL-diaminopimelic acid and the absence of characteristic sugars in the cell wall (Lechevalier and Lechevalier, 1970).




Fig.1: Color of aerial mycelia (White, Grey, Grey), type of spore chains (RF, RF, Spiral) and spore surfaces (Smooth) of three *Streptomyces* isolates IS-01, IS-02, and IS-03, respectively.

Antiviral Activities of *Streptomyces* against TMV:

Recently microbial metabolites have been considered as a potential alternative for chemical insecticides. This is due to the advantages of these metabolites with respect to good environmental compatibility, low mammalian toxicity and a unique mode of action (Seiber 2011). Chen *et al.* (2019) showed that based on the cultural, morphological, physiological and biochemical properties the isolate having antiviral activity against TMV was identified as *Streptomyces ahygroscopicus* and molecularly confirmed by 16S rRNA sequences. Yeo *et al.* (1998) tested the antiviral of an identified isolate of *Streptomyces* coded B25 against TMV under field conditions. They showed that TMV infection decreased from 95.3% up to 58.3% two weeks post-treatment.

Scholthof *et al.* (2011) reported that plant diseases caused by viruses were considered an economic pathogen inducing yearly losses in the production of horticulture and cereal worldwide. They

also showed development and commercialization of highly effective antiviral substances are urgent because chemical insecticides and pesticides were not effective enough to control plant virus disease.

In this study inhibition rates of the three streptomycete isolates against TMV were assessed by the half-leaf method using *N. glutinosa* indicator host as mentioned in the materials and methods. Results in Table (1) and illustrated by Figures (1-5) showed that the strongest anti-TMV effect with an inhibition percentage of 97.5 was recorded for *S. rochei*-IS-02 followed by *S. caeai*-IS-03 (95.5%) and *S. aureocirculatus*-IS-01 (92.5%) at dilution $\frac{1}{2}$, and by increasing the dilutions the inhibitory activities were decreased and reached 37.5% at dilution $\frac{1}{16}$. The experimental results were in harmony with that of Zhu (2011) who found that the fermented filtrate extracted and purified from *Streptomyces parvus* Yn168 inhibited the activity of TMV by reduction of the number of NLL in

which the percentage of inhibition reached up to 84.2% compared to control.

Also, the experimental results could be supported by some investigations, *i.e.*, Han *et al.* (2014) showed that a number of bio-agents isolated from the fermentation broth of *Streptomyces noursei* var. *xichangensis* were able to inhibit assembly of virions among inhibiting the polymerization of TMV coat protein (CP) and induced pathogenesis-

related proteins (PRs), and Han *et al.* (2015) who purified a novel glycoprotein (GP-1) from *Streptomyces kanasensis* exhibited extensive inhibitory effect on TMV infection. In addition, Han *et al.* (2021) showed that the bioactive glycoprotein compound named GP-1 enhanced systemic resistance with a significant reduction in NLL induced by TMV on the leaves of the indicator host.

Table 1: Antiviral activities against TMV of the three *Streptomyces* isolates on *N. glutinosa* indicator host.

<i>Streptomyces</i> isolates	No. of NLL and % of inhibition against TMV							
	1/2		1/4		1/8		1/16	
	NL L	% Inhibition	NLL	% Inhibition	NLL	% Inhibition	NLL	% Inhibition
<i>S. aureocirculatus</i> -IS-01	3	92.5	6	85.0	9	77.5	25	37.5
<i>S. rochei</i> -IS-02	1	97.5	4	90.0	9	77.5	18	55.0
<i>S. caeaoi</i> -IS-03	2	95.5	5	87.5	9	77.5	21	47.5
Control (untreated)	40							




Fig.2: Antiviral activity of *S. aureocirculatus*-IS-01 against TMV on *N. glutinosa* indicator host.




Fig. 3: Antiviral activity of *S. rochei*-IS-02 against TMV on *N. glutinosa* indicator host.




Fig. 4: Antiviral activity of *S. caeai*-IS-03 against TMV on *N. glutinosa* indicator host.



Fig.5: *N. glutinosa* indicator host inoculated with TMV as a control (untreated).

16SrRNA Gene Sequencing and Analysis:

16SrRNA gene was considered a gold taxonomy tool based on the determination of the phylogenetic relationship between the bacteria. In the case of actinobacteria, the large size of *Streptomyces* members makes this tool insufficient to distinguish the closely related species of *Streptomyces* (Law *et al.* (2018). Based on these findings, the need for an alternative phylogenetic gene marker plus 16SRNA gene is an important issue such as RAPD, ISSR, SSR and SCoT methods. DNA fingerprinting of streptomycetes was studied *via* different molecular tools including RAPD-PCR (Mahfouz and Mohamed, 2002; Abdel-Fattah, 2005; Saleh *et al.*, 2011; Shash, 2011; Mohamed *et al.*, 2012; Shori *et al.*, 2012; Mohamed *et al.*, 2013).

In this study, data in Figures (6), (7) and (8) show the numbers 771, 663 and 693 nts represent the partial nucleotide sequences of the 16SrRNA genes of the three streptomycetes, i.e., *S. aureocirculatus*-IS-01, *S. rochei*-IS-02 and *S. caeaoi*-IS-03, of this study, respectively. These isolates were documented in GenBank as strains under the accession numbers LC784317.1, LC784318.1 and LC784319.1, respectively. Results of sequences producing significant alignments showed that the query cover percent between the partial nucleotide sequences of the three *Streptomyces* species and the most five similar strains in GenBank ranged from 96 to 100 for *S. aureocirculatus*-IS-01 (Table 2), 99% for *S. rochei*-IS-02 (Table 3) and 100 for *S. caeaoi*-IS-03 (Table 4).

```

1 ttagtggcga acgggtgagt aacacgtggg caatctgccc tgcaactctgg gacaagccct
61 ggaaacgggg tctaataaccg gataaactct ctgctctcct gagcagaggt tgaaagctcc
121 ggcggtgcag gatgagcccg cggcctatca gcttgttggg gaggtaatgg ctaccaag
181 cgacgacggg tagccggcct gagagggcga cgggccacac tgggactgag acacggccca
241 gactcctacg ggaggcagca gtggggaata ttgcacaatg ggcgaaagcc tgatgcagcg
301 acgccgcgtg agggatgacg gccttcgggt tgtaaacctc tttcagcagg gaagaagcga
361 aagtgcagggt acctgcagaa gaagcgcggg ctaactacgt gccagcagcc gcgtaatac
421 gtagggcgca agcgttgtcc ggaattatg ggcgtaaaga gctcgtaggc ggcttgtcac
481 gtcggttgtg aaagcccggg gcttaacccc gggctctgag tcgatacggg cgagctagag
541 tgtggtaggg gagatcggaa ttcttggtgt agcggtgaaa tgcgcagata tcaggaggaa
601 caccggggc gaagggcgat ctctggcca atactgacgc tgaggagcga aagcgtggg
661 agcgaacagg attagatacc ctggtagtcc acgccgtaaa cgttggggc taggtgtggg
721 caacattcca cgttgtccgt gccgcagcta acgcattaag tgccccgct g

```

Fig. 6: Partial nucleotide sequences (771 nts) of 16SrRNA gene of *Streptomyces aureocirculatus*-IS-01.

```

1 gtcgaaagct ccggcggtgc aggatgagcc cggcgccat cagctagttg gtgaggtaat
61 ggctcaccaa ggcgacgacg ggtagccggc ctgagagggc gaccggccac actgggactg
121 agacacggcc cagactccta cgggagggcag cagtggggaa tattgcacaa tgggcaaaag
181 cctgatgcag cgacgccgcg tgagggatga cggccttcgg gttgtaaac tctttcagca
241 gggaagaagc gaaagtgcag gtacctgcag aagaagcggc ggctaactac gtgccagcag
301 ccgcggtaat acgtagggcg caagcgttgt ccggaattat tgggcgtaaa gagctcgtag
361 gcggcttgtc acgtcgggtg tgaaagccc gggcctaacc ccgggtctgc agtcgatacg
421 ggcaggctag agttcggtag gggagatcgg aattcctggt gtagcgggtg aatgcgcaga
481 tatcaggagg aacaccggtg gcgaaggcgg atctctgggc cgatactgac gctgaggagc
541 gaaagcgtgg ggagcgaaca ggattagata ccctggtagt ccacgccgta aacgggtggc
601 actaggtgtg ggcaacattc cacgttgtcc gtgccgcagc taacgcatta agtgccccgc
661 atg

```

Fig. 7: Partial nucleotide sequences (663 nts) of 16SrRNA gene of *Streptomyces rochei*-IS-02.

```

1 cgttgggctc taggtgtggg cgacattcca cgttgtccgt gccgcagcta acgcattaag
61 tgccccgcct ggggagtagc gccgcaaggc taaaactcaa aggaattgac gggggcccgc
121 acaagcggcg gagcatgtgg ctttaattcga cgcaacgcga agaaccttac caaggcttga
181 catacaccgg aaaactctgg agacagggtc cccctttggg tcgggtgtaca ggtgtgcat
241 ggtgtcgtc agctcgtgtc gtgagatggt ggggtaagtc ccgcaacgag cgcaaccctt
301 atcctgtggt gccagcatgc ttttcggggt gttggggact cacgggagac cgccggggtc
361 aactcggagg aaggtgggga cgacgtcaag tcatcatgcc ccttatgtct tgggctgcac
421 acgtgctaca atggccggta caatgagctg cgataccgcg aggtggagcg aatctcaaaa
481 agcgggtctc agttcggatt ggggtctgca actcgacccc atgaagtcgg agtcgctagt
541 aatcgcagat cagcattgct gcgggtgaata cgttccccgg ccttgtacac accgcccgtc
601 agtcacgaa agtcggtaac accggaagcc ggtggcccaa ccccttgtgg gaggagctg
661 tcgaaggtgg gactggcgat tgggacgaag tcg

```

Fig. 8: Partial nucleotide sequences (693 nts) of 16SrRNA gene of *Streptomyces caeai*-IS-03.

Table 2: Identities (%) and Query Cover (%) with E-value (0%) of partial nucleotide sequences of 16SrRNA gene of *Streptomyces aureocirculatus*-IS-01 compared to the five most similar strains documented in GenBank.

Description	Query Cover (%)	Identities (%)	Accession
<i>Streptomyces aureocirculatus</i> strain DSM 40386T 16S ribosomal RNA gene, partial sequence	100	98.56	KF772674.1
<i>Streptomyces aureocirculatus</i> strain NBRC 13018 16S ribosomal RNA, partial sequence	100	98.56	NR_112344.1
<i>Streptomyces aureocirculatus</i> strain CSSP728 16S ribosomal RNA, partial sequence	98	98.54	NR_043371.1
<i>Streptomyces aureocirculatus</i> strain P6F55 16S ribosomal RNA gene, partial sequence	96	98.51	MN421297.1
<i>Streptomyces aureocirculatus</i> strain YQX2-5m 16S ribosomal RNA gene, partial sequence	96	98.50	OM534621.1

Table 3: Identities (%) and Query Cover (%) with E-value (0%) of partial nucleotide sequences of 16SrRNA gene of *Streptomyces rochei*-IS-02 compared to the five most similar strains documented in GenBank.

Description	Query Cover (%)	Identities (%)	Accession
<i>Streptomyces rochei</i> strain PU-KB10-5 16S ribosomal RNA gene, partial sequence	99	100.00	MN813140.1
<i>Streptomyces rochei</i> strain NEAE-25 16S ribosomal RNA gene, partial sequence	99	100.00	HQ889312.2
<i>Streptomyces rochei</i> strain WR15 16S ribosomal RNA gene, partial sequence	99	99.85	MT659661.1
<i>Streptomyces rochei</i> strain SA1019 16S ribosomal RNA gene, partial sequence	99	99.85	MT355863.1
<i>Streptomyces rochei</i> ANH gene for 16S ribosomal RNA, partial sequence	99	99.85	LC537844.1

Table 4: Identities (%) and Query Cover (%) with E-value (0%) of partial nucleotide sequences of 16SrRNA gene of *Streptomyces caeaoi*-IS-03 compared to the five most similar strains documented in GenBank.

Description	Query Cover (%)	Identities (%)	Accession
<i>Streptomyces caeaoi</i> gene for 16S rRNA, partial sequence, strain: NBRC 12837	100	98.56	AB184183.1
<i>Streptomyces caeaoi</i> strain DSD2595 16S ribosomal RNA gene, partial sequence	100	98.56	MW217135.1
<i>Streptomyces caeaoi</i> strain NBRC 12748 16S ribosomal RNA, partial sequence	100	98.56	NR_041061.1
<i>Streptomyces caeaoi</i> strain SCSIO 68063 16S ribosomal RNA gene, partial sequence	100	98.41	OP482267.1
<i>Streptomyces caeaoi</i> strain Z1-1 16S ribosomal RNA gene, partial sequence	100	98.41	KJ571100.1

The identities between reflect the query cover percent as it ranged from 98.50 to 98.56% between the sequences of *S. aureocirculatus*-IS-01 and those in GenBank represented by accession numbers of KF772674.1, NR_112344.1, NR_043371.1, MN421297.1 and OM534621.1. In the case of *S. rochei*-IS-02 the identities ranged from 99.85 to 100% Between MN813140.1, HQ889312.2, MT659661.1, MT355863.1 and LC537844.1. Regarding *S. caeaoi*-IS-03

the identities compared to AB184183.1, MW217135.1, NR_041061.1, OP482267.1 and KJ571100.1 ranged from 98.41-98.56.

The resulting dendrograms shown in Figures (9), (10) and (11) showed the genetic relationship of partial nucleotide sequences of 16SrRNA gene of the three isolates and that of the five most similar strains documented in GenBank. Results confirmed that these isolates could be strains of *S. aureocirculatus*, *S. rochei* and 100 for *S. caeaoi*.




Fig. 9: Dendrogram shows the genetic relationship of partial nucleotide sequences of 16SrRNA gene of *Streptomyces aureocirculatus*-IS-01 compared to the five most similar strains documented in GenBank.




Fig. 10: Dendrogram shows the genetic relationship of partial nucleotide sequences of 16SrRNA gene of *Streptomyces rochei*-IS-02 compared to the five most similar strains documented in GenBank.




Fig. 11: Dendrogram shows the genetic relationship of partial nucleotide sequences of 16SrRNA gene of *Streptomyces caeai*-IS-03 compared to the five most similar strains documented in GenBank.

RAPD DNA fingerprinting

RAPD-PCR technique allowed the rapid, sensitive and specific evaluation of genetic diversity among *Streptomyces* species (Martin *et al.* 2000). Gharaibeh *et al.* (2003) used the RAPD-PCR technique to evaluate the relatedness of 73 *Streptomyces* isolates producing antibiotics isolated from different soil habitats in Jordan. A number of three monomorphic DNA fragments with sizes of 2777, 800 and 250 pb were common f In 95% of

streptomycete isolates. They confirmed that DNA fingerprinting produced via RAPD-PCR successfully differentiated between the isolates which were characterized with different cultural properties.

In the present study, data in Tables (5), (6), (7) and (8) show the scoring of DNA polymorphisms produced by the applied eight oligonucleotides (P-01, P-02, P-03, P-04, P-05, P-06, P-07 and P-08) used for RAPD-PCR to determine the genetic variation between the three *Streptomyces*

(*S. aureocirculatus*-IS-01, *S. rochei*-IS-02 and *S. caeaoi*-IS-03) of this study. These primers produced a total amplified fragments of 85 distributed as follows 8, 10, 13, 12, 12, 10, 11 and 9 for the eight oligonucleotide primers, respectively (Table 9 and Figs. 12-15). The three *Streptomyces* species appeared 48, 42 and 42 DNA fragments representing percentages of 56.47, 49.41 and 49.41 when the DNA extracts of *S. aureocirculatus*-IS-01, *S. rochei*-IS-02 and *S. caeaoi*-IS-03, respectively, were used as templates in RAPD-PCR test.

Results in Table (10) revealed the type of the 85 amplified DNA fragments, which were classified as four monomorphic fragments and 81 polymorphic fragments. These fragments contained unique DNA fragments as positive DNA markers distributed between the three *Streptomyces*

species as follows: 16 (19.75%), 18 (22.22%) and 14 (17.26%), respectively.

The similarities between the three *Streptomyces* species based on RAPD-PCR DNA fingerprinting were 76.158 between *S. aureocirculatus*-IS-01 and *S. rochei*-IS-02; 66.332 between *S. aureocirculatus*-IS-01 and *S. caeaoi*-IS-03; 73.485 between *S. rochei*-IS-02 and *S. caeaoi*-IS-03 (Table 11). This was confirmed by the phylogenetic relationships in Figure (16) as *S. aureocirculatus*-IS-01 and *S. rochei*-IS-2 were found together in the same cluster, while, *S. caeaoi*-IS-03 was found in a separate cluster. The experimental results were in harmony with that reported by Mahfouz and Mohamed (2002), Hussein et al. (2006), Luo et al. (2010), Mohamed et al. (2012), Shori et al. (2012), Mohamed et al. (2013), Maleki et al. (2013), Biswas et al. (2017), Qattan and Khattab (2019) and Shoukry et al. (2019).

Table 5: Scoring of DNA polymorphisms of three bacterial isolates amplified using P-01 and P-02 primers among ISSR method.

DNA Fragments	Bacterial isolates		
	<i>S. aureocirculatus</i> -IS-01	<i>S. rochei</i> -IS-02	<i>S. caeaoi</i> -IS-03
P-0101	0	0	1
P-0102	1	0	0
P-0103	0	0	1
P-0104	0	0	1
P-0105	1	0	1
P-0106	1	0	0
P-0107	1	0	1
P-0108	0	1	0
Total	4	1	5
P-0201	0	1	0
P-0202	0	1	1
P-0203	1	0	0
P-0204	0	1	1
P-0205	1	0	0
P-0206	0	1	0
P-0207	1	0	0
P-0208	1	0	1
P-0209	0	1	0
P-0210	0	1	0
Total	4	6	3




Fig. 12: Agarose gels (1.2%) stained with ethidium bromide show DNA polymorphisms of three bacterial isolates amplified using P-01 and P-02 primers among ISSR method.

Table 6: Scoring of DNA polymorphisms of three bacterial isolates amplified using P-03 and P-04 primers among ISSR method.

DNA Fragments	Bacterial isolates		
	<i>S. aureocirculatus</i> -IS-01	<i>S. rochei</i> -IS-02	<i>S. caeaii</i> -IS-03
P-0301	1	0	0
P-0302	1	1	0
P-0303	0	1	0
P-0304	0	0	1
P-0305	1	1	1
P-0306	1	0	1
P-0307	1	0	1
P-0308	0	1	0
P-0309	0	1	1
P-0310	1	1	1
P-0311	1	1	0
P-0312	1	1	1
P-0313	0	1	0
Total	8	9	7
P-0401	0	1	0
P-0402	0	0	1
P-0403	1	0	1
P-0404	1	0	0
P-0405	1	0	1
P-0406	0	0	1
P-0407	0	1	0
P-0408	0	1	0
P-0409	0	1	0
P-0410	1	0	1
P-0411	1	1	0
P-0412	1	1	0
Total	6	6	5




Fig.13: Agarose gels (1.2%) stained with ethidium bromide show DNA polymorphisms of three bacterial isolates amplified using P-03 and P-04 primers among ISSR method.

Table 7: Scoring of DNA polymorphisms of three bacterial isolates amplified using P-05 and P-06 primers among ISSR method.

DNA Fragments	Bacterial isolates		
	<i>S. aureocirculatus</i> -IS-01	<i>S. rochei</i> -IS-02	<i>S. caeaoi</i> -IS-03
P-0501	1	0	0
P-0502	0	1	1
P-0503	0	1	1
P-0504	1	0	0
P-0505	0	1	0
P-0506	0	1	1
P-0507	0	0	1
P-0508	0	1	0
P-0509	1	0	1
P-0510	0	0	1
P-0511	1	0	0
P-0512	1	0	0
Total	5	5	6
P-0601	1	0	1
P-0602	1	0	1
P-0603	1	0	1
P-0604	1	1	0
P-0605	1	1	0
P-0606	0	1	0
P-0607	0	1	0
P-0608	1	1	0
P-0609	1	0	0
P-0610	1	0	1
Total	8	5	4




Fig.14: Agarose gels (1.2%) stained with ethidium bromide show DNA polymorphisms of three bacterial isolates amplified using P-05 and P-06 primers among ISSR method.

Table 8: Scoring of DNA polymorphisms of three bacterial isolates amplified using P-07 and P-08 primers among ISSR method.

DNA Fragments	Bacterial isolates		
	<i>S. aureocirculatus</i> -IS-01	<i>S. rochei</i> -IS-02	<i>S. caeaii</i> -IS-03
P-0701	0	0	1
P-0702	1	1	1
P-0703	1	0	1
P-0704	1	0	0
P-0705	1	1	1
P-0706	0	1	1
P-0707	0	0	1
P-0708	1	1	0
P-0709	0	0	1
P-0710	1	0	0
P-0711	0	1	0
Total	6	5	7
P-0801	0	1	1
P-0802	1	0	0
P-0803	1	0	0
P-0804	1	1	1
P-0805	1	0	1
P-0806	1	0	1
P-0807	1	1	1
P-0808	1	1	0
P-0809	0	1	0
Total	7	5	5




Fig. 15: Agarose gels (1.2%) stained with ethidium bromide show DNA polymorphisms of three bacterial isolates amplified using P-07 and P-08 primers among ISSR method.

Table 9: Summary of scoring of DNA polymorphisms of three bacterial isolates amplified using eight primers among ISSR method.

DNA Fragments	TAFs	Bacterial isolates		
		<i>S. aureocirculatus</i> -IS-01	<i>S. rochei</i> -IS-02	<i>S. caeaoi</i> -IS-03
P-01	08	4	1	5
P-02	10	4	6	3
P-03	13	8	9	7
P-04	12	6	6	5
P-05	12	5	5	6
P-06	10	8	5	4
P-07	11	6	5	7
P-08	09	7	5	5
Total	85	48	42	42
%	100	56.47	49.41	49.41

TAFs: Total amplified fragments.


Table 10: DNA unique DNA fragments (Markers) of three bacterial isolates amplified using eight primers among ISSR method.

DNA Fragments	TAFs	MMFs	PMFs	Bacterial isolates		
				<i>S. aureocirculatus</i> -IS-01	<i>S. rochei</i> -IS-02	<i>S. caeaoi</i> -IS-03
P-01	08	0	8	2	1	3
P-02	10	0	10	3	4	0
P-03	13	1	12	1	3	1
P-04	12	0	12	1	4	2
P-05	12	0	12	4	2	2
P-06	10	0	10	1	2	0
P-07	11	1	10	2	1	3
P-08	09	2	7	2	1	3
Total	85	4	81	16	18	14
%	100	4.71	95.29	19.75	22.22	17.28

TAFs: Total amplified fragments. PMFs: Polymorphic fragments. MMFs: Monomorphic fragments.

Table 11: Similarities between the three bacterial isolates based on analysis of DNA polymorphisms produced *via* ISSR method using eight primers.

DNA Fragments	<i>S. aureocirculatus</i> -IS-01	<i>S. rochei</i> -IS-02	<i>S. caeaoi</i> -IS-03
<i>S. aureocirculatus</i> -IS-01	100		
<i>S. rochei</i> -IS-02	76.158	100	
<i>S. caeaoi</i> -IS-03	66.332	73.485	100

**Fig. 16:** Dendrogram shows the genetic relationship between the three bacterial isolates based on analysis of DNA polymorphisms produced *via* ISSR method using eight primers.**Declarations:****Ethical Approval:** It is not applicable.**Conflicts of Interest:** The authors declare that they have no conflict of interest.**Authors Contributions:** I hereby verify that all authors mentioned on the title page have made substantial contributions to the conception and design of the study, have thoroughly reviewed the manuscript, confirm the accuracy and authenticity of the data and its interpretation, and consent to its submission.**Funding:** No funding was received.**Availability of Data and Materials:** All datasets analysed and described during the present study are available from the corresponding author upon reasonable request.**Acknowledgements:** Not applicable.**REFERENCES**

Abd El-Aziz, N.M.; Khalil, B.E. and Ibrahim, H.F. (2023). Enhancement of feather degrading keratinase of *Streptomyces swerraensis* KN23, applying mutagenesis and statistical

optimization to improve keratinase activity. *BMC Microbiology*, 23(1): 158.

Abdel-Fattah, H.I. (2005). Cultural, morphological, physiological and molecular studies on some streptomycete isolates. *Arab Universities Journal of Agricultural Sciences*, 13(2) : 249-268.

Alvarez-Mico, X.; Jensen, P.R.; Fenical, W. and Hughes, C.C. (2013). Chlorizidine, a Cytotoxic 5 H-Pyrrolo [2, 1-a] isoindol-5-one-Containing Alkaloid from a Marine *Streptomyces* sp. *Organic Letters*, 15(5): 988-991.

Anderson, A.S. and Wellington, E.M. (2001). The taxonomy of *Streptomyces* and related genera. *International Journal of Systematic and Evolutionary Microbiology*, 51(3): 797-814.

Biswas, K.; Choudhury, J.D.; Mahansaria, R.; Saha, M. and Mukherjee, J. (2017). *Streptomyces euryhalinus*

- sp. nov., a new actinomycete isolated from a mangrove forest. *The Journal of Antibiotics*, 70(6): 747-753.
- Bo, A.B.; Kim, J.D.; Kim, Y.S.; Sin, H.T.; Kim, H.J.; Khaitov, B.; Ko, Y.K.; Park, K.W. and Choi, J.S. (2019). Isolation, identification and characterization of *Streptomyces* metabolites as a potential bioherbicide. *PLoS One*, 14(9): e0222933.
- Bornet, B. and Branchard, M. (2001). Nonanchored inter simple sequence repeat (ISSR) markers: reproducible and specific tools for genome fingerprinting. *Plant Molecular Biology Reporter*, 19: 209-215.
- Bornet, B.; Muller, C.; Paulus, F. and Branchard, M. (2002). Highly informative nature of inter simple sequence repeat (ISSR) sequences amplified using tri- and tetra-nucleotide primers from DNA of cauliflower (*Brassica oleracea* var. botrytis L.). *Genome*, 45(5): 890-896.
- Chen, J.; Liu, H.; Xia, Z.; Zhao, X.; Wu, Y. and An, M. (2019). Purification and structural analysis of the effective anti-TMV compound ϵ -poly-L-lysine produced by *Streptomyces ahygroscopicus*. *Molecules*, 24(6):1156.
- El-Ahdal, A. M. E.; Mayer, F.; El Afifi, S. I. and Allam, E. K. (1984). Characterization of two strains of Tobacco mosaic Virus by biological, biochemical, immunological and electron microscopical techniques. *Zeitschrift für allgemeine Mikrobiologie*, 24(8): 507-514.
- El-Masry, S.S.; Sadik, M.S. and Akl, B.A. (2019). Efficiency of two molecular tools based on dna used for differentiating some microbial strains. *Arab Universities Journal of Agricultural Sciences*, 27(1): 219-229.
- El-Masry, S.S.; El-DougDoug, K.A.; Othman, B.A. and Mohamed, S.H. (2021). Isolation and Characterization of Antagonistic Activity of Some Streptomyces and their Specific Actinophages from Soil. *Pakistan Journal of Biological Sciences: PJBS*, 24(4): 516-526.
- Fedriago, K.; Giacomini, R.M.; Faria, C.M. and Da-Silva, P.R. (2016). ISSR primers for analysis of genetic variability of *Stenocarpella maydis*. *Tropical Plant Pathology*, 41: 270-275.
- Fraile, A., and García-Arenal, F. (2018). Tobamoviruses as models for the study of virus evolution. *Advances in virus research*, 102: 89-117.
- Gharaibeh, R.; Saadoun, I. and Mahasneh, A. (2003). Genotypic and phenotypic characteristics of antibiotic-producing soil *Streptomyces* investigated by RAPD-PCR. *Journal of Basic Microbiology*, 43(1): 18-27.
- Gooding, G. V. and Hebert, T. T. (1967). A simple technique for purification of tobacco mosaic virus in large quantities. *Phytopathology*, 57 (11): 1285-1285.
- Han, L.; Zhang, G.; Miao, G.; Zhang, X. and Feng, J. (2015). *Streptomyces kanasensis* sp. nov., an antiviral glycoprotein producing actinomycete isolated from forest soil around Kanas Lake of China. *Current microbiology*, 71: 627-631.
- Han, L.; Sun, Y.; Zhou, X.; Hao, X.; Wu, M.; Zhang, X. and Feng, J. (2021). A novel glycoprotein from *Streptomyces* sp. triggers early responses of plant defense. *Pesticide Biochemistry and Physiology*, 171: 104719.
- Han, Y.; Luo, Y.; Qin, S.; Xi, L.; Wan, B. and Du, L. (2014). Induction of

- systemic resistance against tobacco mosaic virus by Ningnanmycin in tobacco. *Pesticide biochemistry and physiology*, 111:14-18.
- Hassanien, H.A.; Elnady, M.; Obeida, A. and Itriby, H. (2004). Genetic diversity of Nile tilapia populations revealed by randomly amplified polymorphic DNA (RAPD). *Aquaculture Research*, 35(6): 587-593.
- Hussein, S.H.; Abdel-Fattah, H.I. and Chaudhry, Z. (2006). A comparative study between a numerical and rapd-pcr methods for the identification of some streptomycete strains. *Pakistan Journal of Biotechnology*, 3(1-2): 71-82.
- Kamjam, M.; Nopnakorn, P.; Zhang, L.; Peng, F.; Deng, Z. and Hong, K. (2019). *Streptomyces polaris* sp. nov. and *Streptomyces septentrionalis* sp. nov., isolated from frozen soil. *Antonie Van Leeuwenhoek*, 112: 375-387.
- Kim, D.; Chun, J.; Sahin, N.; Hah, Y.C. and Goodfellow, M. (1996). Analysis of thermophilic clades within the genus *Streptomyces* by 16S ribosomal DNA sequence comparisons. *International Journal of Systematic and Evolutionary Microbiology*, 46(2): 581-587.
- Kim, B.; Sahin, N.; Minnikin, D.E.; Zakrzewska-Czerwinska, J.; Mordarski, M. and Goodfellow, M. (1999). Classification of thermophilic streptomycetes, including the description of *Streptomyces thermoalcalitolerans* sp. nov. *International Journal of Systematic and Evolutionary Microbiology*, 49(1): 7-17.
- Korn-Wendisch, F. and Kutzner, H.J. (1992). The family Streptomycetaceae. *Prokaryotes*, (Ed. 2), pp.921-995.
- Law, J.W.F.; Tan, K.X.; Wong, S.H.; Ab Mutalib, N.S. and Lee, L.H. (2018). Taxonomic and characterization methods of *Streptomyces*: a review. *Progress In Microbes & Molecular Biology*, 1(1).
- Lechevalier, M.P. and Lechevalier, H. (1970). Chemical composition as a criterion in the classification of aerobic actinomycetes. *International Journal of Systematic and Evolutionary Microbiology*, 20(4): 435-443.
- Luo, J.; Li, J.; Liu, D.; Zheng, Y.; Liu, Y. and Wang, M. (2010). Application of RAPD assays in analyzing streptomycetes *gilvosporeus* strains from genome shuffling. In *2010 4th International Conference on Bioinformatics and Biomedical Engineering* (pp. 1-4). IEEE.
- Mahfouz, H.T. and Mohamed, S.H. (2002). Physiological antagonistic and fingerprinting studies on some halotolerant *Streptomyces* strains. *Arab Journal of Biotechnology*, 5(1): 103-120.
- Maleki, H.; Dehnad, A.; Hanifian, S. and Khani, S. (2013). Isolation and molecular identification of *Streptomyces* spp. with antibacterial activity from northwest of Iran. *BioImpacts: BI*, 3(3):129.
- Malkawi, H.I.; Saadoun, I.; Moumani, F.A. and Meqdam, M.M. (1999). Use of RAPD-PCR fingerprinting to detect genetic diversity of soil *Streptomyces* isolates. *The New Microbiologica*, 22(1): 53-58.
- Martin, P.; Dary, A.; André, A. and Decaris, B. (2000). Identification and typing of *Streptomyces* strains: evaluation of interspecific, intraspecific and intraclonal differences by RAPD fingerprinting. *Research in Microbiology*, 151(10):853-864.

- Mehling, A.; Wehmeier, U.F. and Piepersberg, W. (1995). Nucleotide sequences of streptomycete 16S ribosomal DNA: towards a specific identification system for streptomycetes using PCR. *Microbiology*, 141(9): 2139-2147.
- Mohamed, S.H.; Abdel-Fattah, H.I.; Selim, S.M. and Sharaf, M.S. (2001). Identification and molecular studies on some halotolerant streptomycetes isolated from Sinai sandy soil. *Arab Journal of Biotechnology*, 4(2) :179-196.
- Mohamed S.H.; Omran, W.M.; Shima, M.A.S.; Al-Shehri, A.S. and Sadik, A.S. (2012). Isolation and Identification of some halotolerant Actinomycetes having antagonistic activities against some plant pathogens (ie, Tobacco mosaic virus, Aspergillus Sp., Fusarium Sp.) from soil of Taif Governorate KSA. *Pakistan Journal of Biotechnology*, 9(1): 1-12.
- Mohamed, S.H.; Al-Saedi, T.A. and Sadik, A.S. (2013). Halotolerant streptomycetes isolated from soil at Taif region, Kingdom of Saudi Arabia II: RAPD-PCR analysis and salt tolerance-gene isolation. *African Journal of Biotechnology*, 12(13).
- Qattan, S.Y. and Khattab, A.A. (2019). Molecular Characterization of Streptomyces albogriseolus Excellent Mutants for Neomycin Production. *Journal of Pure & Applied Microbiology*, 13(3).
- Roossinck, M.J. (2015). Plants, viruses and the environment: Ecology and mutualism. *Virology*, 479: 271-277.
- Saleh, E.A.; Zaki, M.M.; Rahal, A.; Mohamed, S.H.; Sadik, A.S. and Behieldin, A.M. (2011). Molecular characterization of some Basta herbicide tolerant streptomycetes. In *The 26th meeting of Saudi Biological Society, " Climatic Change and Biodiversity" Taif University, Taif.* pp: 7-9.
- Sambrook, J.; Fritsch, E.F. and Maniatis, T. (1989). Gel electrophoresis of DNA In: *Molecular Cloning: A Laboratory Manual, Part 6. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York,* pp:1-15.
- Scholthof, K. B. G.; Adkins, S.; Czosnek, H.; Palukaitis, P.; Jacquot, E.; Hohn, T. Hohn, B.; Saunders, K.; Candresse, T.; Ahlquist, P. and Foster, G. D. (2011). Top 10 plant viruses in molecular plant pathology. *Molecular Plant Pathology*, 12(9): 938-954.
- Seiber, J.N. (2011). Sustainability and agricultural and food chemistry. *Journal of agricultural and food chemistry*, 59(1):1-21.
- Shash, S.M. (2011). Somaclonal variation of three tolerance Streptomyces isolated from saline soil. *New York Science Journal*, 4(12): 67-73.
- Shori, G.B.O.; Mohamed Sonya, H.; Abdel-Salam Shima, M. and Sadik, A.S. (2012). Characterization of streptomycetes having antibiosis activities isolated from soil in western region of KSA. *Pakistan Journal of Biotechnology*, 9(1): 55-60.
- Shoukry, A.A.; Khatab, I.A.; Faried, M.A. and Khattab, A.A. (2019). Isolation, Evaluation and Molecular Identification of Streptomyces Isolates with Antimicrobial Activities. *Journal of Agricultural Chemistry and Biotechnology*, 10(5): 103-112.
- Sneath, P.H. and Sokal, R.R. (1973). *Numerical taxonomy. The principles and practice of numerical classification.* WF

- Freeman & Co., San Francisco, p:573.
- Stackebrandt, E.; Witt, D.; Kemmerling, C.; Kroppenstedt, R. and Liesack, W. (1991). Designation of streptomycete 16S and 23S rRNA-based target regions for oligonucleotide probes. *Applied and Environmental Microbiology*, 57(5): 1468-1477.
- Takeuchi, T.; Sawada, H.; Tanaka, F. and Matsuda, I. (1996). Phylogenetic analysis of *Streptomyces* spp. causing potato scab based on 16S rRNA sequences. *International Journal of Systematic and Evolutionary Microbiology*, 46(2): 476-479.
- Waksman, S.A. and Henrici, A.T. (1943). The nomenclature and classification of the actinomycetes. *Journal of bacteriology*, 46(4): 337-341.
- Williams, S.T. (1989). Genus *Streptomyces* Waksman and Henrici 1943. *Bergey's Manual of Systematic Bacteriology*, 4: 2452-2492.
- Williams, S.T.; Goodfellow, M.; Alderson, G.; Wellington, E.M.H.; Sneath, P.H.A. and Sackin, M.J. (1983). Numerical classification of *Streptomyces* and related genera. *Microbiology*, 129(6): 1743-1813.
- Yeo, W. H.; Kim, Y. H.; Kim, Y. S.; Kim, S. S. and Chae, S. Y. (1998). Control effects of an antibiotic produced by *Streptomyces* sp. B25 on tobacco mosaic virus and determination of its molecular structure. *Journal of the Korean Society of Tobacco Science*, 20(2): 172-177.
- Zhu, H.J. (2011). Anti-TMV activity and active components of metabolites of *Streptomyces* HNS2-2. *Chinese Journal of Biological Control*, 27(4): 515.